

Karle Medical Group, P.C.

455 Barclay Circle, Suite D

Rochester Hills, MI 48307

T: 248-852-9596 | F: 248-852-9453

Christine L. Karle, D.O.
Tracey R. Ticcony, N.P.C.

Bridget C. Karle, M.D.
Malaz Alatassi, M.D.

Kristie Burkland, N.P.C.
Molly Bylsma, N.P.C. Amir Sankari, M.D.

Denise Gavorin, D.O.
Lindsay Runft, D.N.P.

Insulin Dosage Schedule

Printed Patient Name: _____

Patient DOB: _____ Date: _____ MA Initials: _____

Long Acting Insulin Instructions:

Levemir / Lantus

*Before Breakfast: _____ units

*Before Bedtime: _____ units

- Increase or decrease dosage by 2 units every 4 days until fasting blood sugar = 90 – 100, then **STOP CHANGING** the dosage.

Fast Acting Insulin Instructions: (Check blood sugar then take insulin 5 – 10 minutes before you eat)

Novolog / HumaLog / Apidra

Take before meals according to the following scale:

Test Sugar Levels at the following intervals: before breakfast, lunch, dinner and bedtime

	Before Breakfast	Before Lunch	Before Dinner	Before Bed
Sugar Level	_____ + Scale	_____ + Scale	_____ + Scale	Long Acting Insulin
<80				
81-100				
101-150				
151-200				
201-250				
251-300				
301-350				
351-400				
>400				
Generic Dose: (if not checking sugars)				

Target sugar before meals should be 100 – 110.

Target sugar after meals should be less than 140.

For activity, subtract _____ units from above pre-meal scale.

For over-eating add _____ units to the above scale.